ABSTRACTS

Broad-spectrum group treatment for parents bereaved by the violent deaths of their 12- to 28year-old children: a randomized controlled trial

Murphy SA, Johnson C, Cain KC, Das Gupta A, Dimond M, Lohan J, Baugher R. Death Studies 1998; 22(3): 209-35

Violent deaths caused by accidents, homicides and suicides account for 80% of all deaths among youth and young adults in the USA each year. This study examines the efficacy of a 10-week intervention programme offered to 153 parents from between two and seven months after the violent deaths of their 12 to 28-year-old children. The bereaved parents were assessed for mental distress, post-traumatic stress symptoms, delayed loss accommodation, deteriorating physical health and marital strain. The aim of the study - the methodology of which is described in great detail - was to provide the parents with information, skills and support which would help them to come to terms with their loss. It was found that mothers who were most distressed responded best, while fathers showed no immediate benefit. The authors suggest that further research is needed to investigate this lack of response on the part of the fathers and to characterise further those who benefit from similar programmes.

Mothers now childless: structures of the life-world

Talbot K. Omega 1997/8; 36(1): 45-62

The author of this moving article is herself a bereaved mother whose only child died at the age of nine. The death of a child, especially an only child, presents the kind of suffering and challenge to self that creates a need to search for the meaning of life. She describes interviews which she conducted with 10 mothers who had lost their only child, and who could be divided into two groups of five those who were unable to visualise the future without their child and those who had made a conscious decision to survive their loss. Significantly, the latter group were all working as volunteers to help other mothers who had lost an only child, which allowed them to keep the mothering (nurturing) part of their identity. She suggests ways of helping such bereaved mothers, above all the need for sympathetic, non-judgemental counsellors and friends who will listen and understand.

Key elements in a framework for helping grieving children and adolescents.

Corr CA, Corr DM. Illness, Crisis and Loss 1998: 6(2); 142-60

It is not easy for adults to accept the death of a loved one; how much more difficult for children and adolescents! This article describes two approaches through which help can be given in such cases. First, that of the Dougy Center in Portland,

USA, which operates in the belief that every bereaved child or adolescent should have an opportunity to grieve his or her loss in a supportive and understanding environment. Secondly, it describes the philosophy of Sandra Fox, who identified four tasks that confront grieving children and adolescents, and discusses ways in which they can be helped to understand their loss and learn how to cope with it. This thoughtful and useful article ends with 10 guidelines for adults who are involved with young people.

Long-term unresolved grief: applying Bowlby's variants to adult survivors of early parental death

Dilworth JL, Hildreth GL. Omega 1997/8; **36**(2): 148-59.

'Short-term grief resolution in children and adults has received considerable attention,' write the authors of this article, but we still know little about the long-term grief resolution of any group.' Attempting to address this lack, they discuss the correlation between loss of a parent in childhood and possible adult pathology resulting from the bereavement. Their study is an extension of Bowlby's disordered variants of childhood grief, and they suggest that behavioural and thought manifestations in adult survivors of early bereavement indicate unresolved grief. They come to no definite conclusions, but stress the need for further research, using appropriate methodological controls. It is surprising to read that, discussing parental worries about dying, in one study none of the mothers in the control group expressed great anxiety about dying and leaving their children bereaved.

Mourning, ritual and recovery after an airline tragedy.

Katz P, Bartone P. Omega 1997/8; 36(3): 193-200.

The crash of a chartered Army plane in Newfoundland, in 1985, resulted in the deaths of 248 soldiers from one battalion stationed in Fort Campbell, Kentucky, whose families all came from that region. This short article first discusses the value of mourning rituals and then describes how, in this particular instance, they made it possible for the families and colleagues of the dead soldiers to come to terms with their traumatic loss. The rituals which were carried out were numerous and elaborate, and it is believed that they helped the bereaved to accept the deaths, re-affirmed the solidarity of the community and enabled individuals and the community to reestablish a sense of control.

Sheila Hodges and John Bush

CORRECTION

Bereavement Care 1998; 17(2): 32 The heading of the final abstract should not have read 'Sibling bereavement...' but 'Communities of feeling: the culture of bereaved parents. Riches G, Dawson P. Mortality 1966; 1(2): 143-61'.

Volume 17: Spring 1998 - Winter 1998

Key: A = article No.1 = Spring issue Ab = abstract No.2 = Summer issue R = review No.3 = Winter issue

A new model of grief; bereavement and biography (Ab); No 1:16 Attachment across the life cycle (R); No 2: 31

Balk D (Ab); No 2: 32 Broad-spectrum group treatment for parents bereaved by the violent deaths of their 12 to 28-year-old children (Ab); No 3: 48

Collier C (A); No 1: 3-6 Coyle A (Ab); No 1: 16 Crossland C (A); No 2: 19-20 Culture and grief (A); No 1: 7-10 Critical Incident Debriefing (R); No 3: 47

Daniel G (A): No 3: 35-8

Death and bereavement across cultures (R); No 1; 15 Death Customs (R); No 1; 15 Death and loss (R); No 2; 31 Dominica Sister F (R); No 2: 30 Drawing the boundaries (A); No 2; 27

Dying, death and bereavement (R): No 2: 32 Experiences of AIDS-related bereavement among gay men: implications for care (Ab): No 1: 16

Family problems after a bereavement (A); No 3: 35-8 Finlay K (A); No 3: 42-3 Folkman S (Ab); No 1: 16 Frean A (A); No 2: 26 From mourning and melancholia to bereavement and biography; an assessment of Walter's new model of grief (Ab); No 1: 16

Gender differences among parents who experience the death of an infant weighing less than 500 grams at birth (Ab); No 2: 32

Grief and isolation after an AIDS-related death (A): No 3: 38-41

Heavenly Hurts (R); No 1: 15 Hébert M (Ab); No 2: 32 Hemmings P (A); No 1: 14 Hemmings P (A); No 2: 27

Jacoby Klein S (R); No 1: 15 Just my reflection (R); No 2: 30 Journey to 'Mum's last mountain' (A); No 1: 3-6

Kavanaugh K (Ab); No 2: 32 Kershaw H (A); No 3: 44 Key elements in a framework for helping grieving children and adolescents (Ab);

Laungani P (R); No 1: 15 Leaman O (R); No 2: 31 Letting go and keeping hold: a reply to Stroebe (Ab); No 1: 16 The lone twin (R); No 2: 31 Long-term unresolved grief: applying Bowlby's variants to adult survivors of early parental death (Ab); No 3: 48

Mahon M (Ab); No 2: 32 Marcus J (R); No 1: 14 Marris P (R); No 2: 31 Maternal responses to the life and death of a child with a developmental disability: a story of hope (Ab); No 1: 16 Merrington B (R); No 2: 30 Milo E (Ab); No 1: 16 Mind mapping with bereaved adolescents (A); No 3: 44 Mothers now childless; structures of the life-world (Ab); No 3: 48

Moss M (Ab); No 1: 16 Moss S (Ab); No 1: 16 Mourning, ritual and recovery after an airline tragedy (Ab); No 3: 48 Munday J (R); No 2: 30

New directions in bereavement research: exploration of gender differences (Ab); No 2: 32

No chance to say goodbye (R); No 1: 14 The Norwegian Bereavement Project (A); No 2: 24-25

O'Regan M (A); No 3; 38-41 Panufnik CJ (R); No 3; 45-6 Parents' support group in Dunblane (A); No 2: 26 Parkes C (R); No 1: 15 Parkes C (A); No 2: 21-23 Parkes C (R); No 2: 31

Perinatal bereavement in its cultural context (Ab); No 2: 32 Psychological development during four

Parkinson F (R): No 3: 48

years of bereavement: a longitudinal case study (Ab); No 2: 32 Reaction of schools in the Bath area to

Reaction of schools in the Bath area to bereavement (A); No 3: 42-3 Resch N (Ab); No 1: 16 Returning to employment after bereave ment (A); No 1: 11-13 Richards T (Ab); No 1: 16 Robinson L (Ab); No 2: 32 The role of gender in middle-aged children's responses to parent death (Ab); No 1: 16 Rushton L (R); No 1: 15 Russell K (A); No 1: 11-13

Sandvik O (A); No 2: 24-25

Schut H (A); No 1: 7-10 Shared experiences: seven books and a film which may help bereaved children (R); No 3: 45-6

Sibling bereavement: a concept analysis (Ab); No 2: 32 Sites of memory, sites of mourning (R); No

2: 31 Social support 'Internet-works', caskets for sale, and more: thanatology and the

information superhighway (Ab); No 1: 16 Sofka CJ (Ab); No 1: 16

Spiritual aspects of loss at time of a partner's death from AIDS (Ab); No 1: 16 St George's Medical School (R); No 2: 32 Stevenson-Hinde J (R); No 2 31

Stevenson-Hinde J (R); No 2 31 Stroebe M (A); No 1: 7-10 Sroebe M (Ab); No 1: 16 Stroebe M (Ab); No 2: 32 Suffering Love (R); No 2: 30

Surviving the death of a child (R): No 2: 30 Thuen F (A): No2: 24-25

Traditional models and theories of grief (A): No 2: 21-23

Trust circles (A); No 1: 14

Vesta L (Ab); No 2: 32

Walter T (Ab); No 1: 16 Winter J (R): No 2: 31 Wohlenhaus-Munday (R); No 2: 30 Woodward J (R); No 2: 31 Working with widowed men (A); No 2: 19 Wright C (Ab); No 1: 16

Young B (R); No 1: 15

